

Running Head: LATENT STRUCTURE OF PSYCHOPATHY

A taxometric analysis of the latent structure of psychopathy: Evidence for dimensionality

Jean-Pierre Guay, Ph.D.

Université of Montréal

&

Institut Philippe-Pinel de Montréal

John Ruscio, Ph.D.

Elizabethtown College

Raymond A. Knight, Ph.D.

Brandeis University

Robert D. Hare, Ph.D.

University of British Columbia

Published in *Journal of Abnormal Psychology*

Copyright 2007 by the American Psychological Association

<http://www.apa.org/journals/abn/>

This article may not exactly replicate the final version published in the APA journal. It is not the copy of record.

Abstract

The taxonomic status of psychopathy is controversial. Whereas some studies have found evidence that psychopathy, at least its antisocial component, is distributed as a taxon, others have found that both major components of psychopathy —callousness/unemotionality and impulsivity/antisocial behavior —appear to distribute as dimensions and show little evidence of taxonicity. In the present study, recent advances in taxometric analysis were added to Meehl's multiple consistency tests strategy for assessing taxonicity (Meehl, 1995), and they were applied to Psychopathy Checklist-Revised (PCL-R) ratings of 4865 offenders sampled from multiple forensic settings. The results indicated that both the individual components of psychopathy and their interface are distributed dimensionally. Both the implications of these results for research in psychopathy and the integration of these findings with previous taxometric studies of psychopathy are discussed.

Key words: Psychopathy, taxometrics, latent structure, antisocial behavior

A taxometric analysis of the latent structure of psychopathy: Evidence for dimensionality

Psychopathy is a personality disorder that comprises a distinct cluster of emotional, interpersonal, and behavioural characteristics (e.g., emotional detachment, callousness, irresponsibility, impulsivity) and that is characterized by a disregard for the societal rules and the rights of others (Hare, 1996). Its association with violence (Porter & Woodworth, 2006) and its usefulness as a risk factor in predicting criminal recidivism (Douglas, Vincent, & Edens, 2006) have increased its prominence in the last decade in both criminology and psychopathology. The origins of our current conceptualization of the construct can be traced to Cleckley's (1976) classic description of the syndrome. He delineated its characteristics without, however, addressing the issue of whether it represents the coalescence of extreme manifestations on a number of dimensional traits or constitutes a taxon, i.e., a distinct, nonarbitrary entity or class.

The first formalized assessment tool for measuring psychopathy was Hare's (1980) operationalization of the construct in the Psychopathy Checklist (PCL) and its revision, the PCL-R (Hare, 1991, 2003). Although Hare conceptualized the PCL as a way of indicating how closely an individual approximated the "prototypic psychopath" and proposed a PCL-R cutting score of 30 (out of 40) to consider a person sufficiently close to the psychopath prototype, he also recognized that viable arguments could be made for using the PCL-R to obtain dimensional scores.

The theoretical case for hypothesizing that psychopathy may be distributed as a taxon revolves around evidence for specific genetic, neurobiological, cognitive, and affective covariates of the construct that are consistent with the hypothesis that a specific etiology or specific etiologies may account for it. Probably the most widely accepted theoretical understanding of taxonicity and the strongest examples of taxa have focused on entities specified

by the conjunction of a distinct pathology and etiology (Meehl, 1973; 1992). Although the establishment of such covariates does not in itself confirm either a syndrome's specific etiology or its taxonicity, and dimensional models can be generated to handle such correlative findings, the *absence* of such covariates would certainly argue against proposing a taxonic distribution.

Two (Harpur, Hakstian, & Hare, 1988; Harpur, Hare, & Hakstian, 1989), three (Cooke & Michie, 2001), and four factor (Hare, 2003; Hare & Neumann, 2006) models of the PCL have all identified two overarching components involving impulsivity-antisocial behavior and affective-interpersonal features. Because these two components have consistently yielded distinct patterns of correlations with characteristics relevant to psychopathy, different models of specific underlying processes have been proposed for each (e.g., Blonigen, Hicks, Krueger, Patrick, & Iacono, 2005; Fowles & Dindo, 2006; Patrick & Zempolich, 1998). Such models raise the possibility that either or both factors may be distributed as taxa. A brief survey of some of the correlative evidence in the genetic, neurobiological, cognitive, and affective domains provides an adequate justification to support a taxonomic investigation.

There is considerable evidence from adoption and twin genetic studies, including those of twins reared apart, that indicate that genetic factors both play a significant role in the likelihood that a person will commit a criminal act (Gottesman & Goldsmith, 1994; Grove, Eckert, Heston, Bouchard, et al., 1990; Mednick, Gabrielli, & Hutchings, 1984) and also increase the probability that an individual will be diagnosed as Antisocial Personality Disorder (APD) during his lifetime (Cadoret, Yates, Troughton, Woodworth, & Stewart, 1995; Ge, Conger, Cadoret, Neiderhiser, Yates, et al., 1996; Lyons et al., 1995). Although behavioral genetic research suggests that the Factor 2 features may have higher heritability than the Factor 1 (Edelbrock, Rende, Plomin, & Thompson, 1995; Krueger, 2000; Mason & Frick, 1994; Depue, 1996, but see Livesley, 1998,

for contrary evidence), there are also data supporting the importance of independent genetic influences in the manifestation of Factor 1 (Taylor, Loney, Bobadilla, Iacono, McGue, 2003; Patrick, 2003).

A number of neurobiological deficits or anomalies have been identified in both psychopathic criminals and “successful” psychopaths, who have largely avoided extensive contact with the criminal justice system or extended incarceration. For instance, both specific neurological structural features in the amygdala, orbito-frontal cortex, and hippocampus (e.g., Blair, 2004; Mitchell, Colledge, Leonard, & Blair, 2002; Raine, 2001; Raine et al., 2004; Tiihonen et al., 2000) and functional anomalies in the amygdala/hippocampal formation, parahippocampal gyrus, ventral striatum, anterior and posterior cingulated gyri, and fronto-temporal cortex (e.g., Kiehl, Hare, Liddle, & McDonald, 1999; Kiehl, Smith, Mendrek, Forster, Hare, & Liddle, 2004; Kiehl, Smith, Hare, Mendrek, Forster, Brink, & Liddle, 2001; Müller et al., 2003; Völlm et al., 2004) have been linked to the syndrome or its components. Some speculations have been proposed about a comprehensive and integrated mapping of these deficits onto psychopathy (Blair, Mitchell, & Blair, 2005) and onto the two major PCL-R factors that assess it (Patrick, 2003). An understanding of the latent distribution of the PCL-R and its factors could substantially advance the search for such an integration.

Psychopaths have been found to be deficient on a number of cognitive (e.g., Hervé, Hayes & Hare, 2001; Morgan & Lilienfeld, 2000; Newman & Lorenz, 2003) and affective tasks (e.g., Blair, 2001; Blair, Mitchell, Richell, Kelly, Leonard, Newman & Scott, 2002; Patrick, 2001; Williamson, Harpur, & Hare, 1991) that map onto the structural features noted above. Psychopaths’ difficulty in shifting a dominant behavior when contingencies have been reversed (Newman & Lorenz, 2003) and their difficulties with working memory and other aspects of

executive functioning (Morgan & Lilienfeld, 2000; Séguin, 2004) implicate deficits in their orbito-frontal cortex. In contrast, psychopaths' impairments in passive avoidance learning (Newman & Kosson, 1986), dysfunctional response to another's sadness or fear (Blair, 1995), reduced augmentation of the startle reflex by threat primes (Levenston, Patrick, Bradley, & Lang, 2000), impaired aversive conditioning (Raine, Venables, & Williams, 1996), and deficient processing of fearful expressions (Blair, Colledge, Murray, & Mitchell, 2001) are perhaps better accounted for at the neural level by deficiencies in the functions of the amygdala (Blair, 2004) or in the integration of frontal-limbic processes (Hare, 2003; Müller et al., 2003). Although speculative models to integrate these deficits have been proposed (e.g., Blair et al., 2005; Fowles & Dindo, 2006), unpacking the complexity of the development of the underlying core processes (Séguin, 2004) and mapping onto the specific behavioral patterns in psychopathy have remained elusive.

Such correlations with specific genetic, neurobiological, cognitive, and affective processes suggest the possibility that specific neurobiological deficiencies may be necessary antecedents of psychopathy. Although dimensional models may be proposed to account for such causes (e.g. Benning, Patrick, Blonigen, Hicks, & Iacono, 2005; Miller, Lynam, Widiger, & Leukefeld, 2001), a taxonic distribution is a viable theoretical possibility that must be investigated. Resolution of this issue of the latent structure of psychopathy is not only important for developing theoretical models of the etiology and course of psychopathy, but it also has critical implications for determining optimal investigative strategies and for specifying the ideal psychometric qualities of scales constructed both for clinical and dispositional decision-making (Krueger, 1999; Meehl, 1992; Ruscio & Ruscio, 2004a). Considering the prominence of PCL-R ratings in risk assessment instruments (e.g., Quinsey, Harris, Rice, & Cormier, 1998), and its

widespread use in legal dispositional decisions (Edens & Petrila, 2006), such scaling issues have substantial practical import. Consequently, a number of studies have addressed the problem of psychopathy's latent distribution (Guay & Knight, 2003; Harris, Rice, & Quinsey, 1994; Marcus, John, & Edens, 2004; Skilling, Quinsey, & Craig, 2001; Vasey, Kotov, Frick, & Loney, 2005), but with mixed results. All of these prior studies unfortunately have suffered from sampling and methodological weaknesses that have limited their ability to provide definitive answers to this question.

The two studies that found no evidence for taxonic latent structures both analyzed self-report data. Guay and Knight (2003) conducted a taxometric investigation of the components of psychopathy using the Multidimensional Assessment of Sex and Aggression (the MASA; Knight & Cerce, 1999). The MASA is a self-report inventory that covers multiple domains (childhood experiences, family and social relationships, school and work experiences, alcohol and drug use, and sexual and aggressive behavior and fantasies) and that was developed to supplement archival records. Participants were 330 sex offenders, 155 generic criminals, and 93 community controls, who had been tested on paper-and-pencil and computerized versions of the MASA. A total of eleven scales measuring conning and superficial charm, emotional detachment and behavioral problems, and impulse control were generated using factor analysis in combination with a Rasch model. The various taxometric techniques (MAMBAC, MAXCOV, MAXEIG) that were assessed all generated results consistent with a dimensional structure, with no evidence of taxonicity. Marcus, John, and Edens (2004) evaluated a sample of 309 incarcerated offenders (51.6% African American, 37.5% Caucasian, 8% Hispanic, 2.9% "other," and 91.3% male) using the Psychopathic Personality Inventory (PPI; Lilienfeld & Andrews, 1996). The authors performed MAMBAC, MAXEIG and L-Mode analyses, and their results showed no evidence of

taxonicity. Both studies may have compromised their ability to identify a taxon, because of the potential for increased nuisance covariation in self-report data that is produced by response styles and biases. Such method variance could have artifactually increased the correlation of scales within the taxon and the complement and thereby reduced the potential to identify a taxon.

Harris, Rice, and Quinsey (1994) analyzed data collected from 653 mentally disordered participants from a maximum-security institution, who had been adjudicated not guilty by reason of insanity. Although the authors argued that their results provided evidence for a taxon, several methodological ambiguities of their study undermine their conclusion. First, their sample comprised a select, potentially biased groups of offenders (Marcus et al., 2004). Second, with the exception of their application of Meehl's taxometric analyses, most of the statistical procedures used by Harris et al. (1994) lacked empirical support as methods for distinguishing between taxonic and dimensional structures. The validity of their iterative methods approach has not been tested by any Monte Carlo studies, and the distributions of Bayesian probabilities can readily be U-shaped even when the latent structure is dimensional, especially when a large number of items are used as in Harris et al.' study. Third, their exclusive use of file reviews introduced severe limitations. The data for accurately rating Factor 1 items in archival files is often missing (Hare, 2003). Moreover, insufficient file information can lead raters to score items for which data are missing or inadequate from information on related items (Alpert, Shaw, Pouget, & Lim, 2002), thereby decreasing item covariation within the putative taxon, increasing item correlation across the taxon and complement, and increasing the potential to find a pseudo-taxon. File reports often focus on salient, egregious Factor 2 antisocial behaviors that may make ratings more vulnerable to raters' a priori taxonic (Beauchaine & Waters, 2003) or item contingency (Bolt, Hare, Vitale, & Newman, 2004) biases, and thereby may increase the probability of pseudo-taxonicity. In this

regard, it is interesting that Harris et al., (1994) only found evidence for a taxon in their Factor 2 analyses. Fourth, the authors used an admixed sample in which they employed some matching procedures. The alleged taxon might have been an artifact of their sampling strategy. This possibility could have been avoided by re-running their analyses within samples, or they could have presented a contingency table of sample membership and taxon/complement assignment. Fifth, Harris et al. (1994) did not have access to the simulation programs used in the present study, which generate taxonic and dimensional comparison data to analyze as an interpretive aid. Without this, they did not notice the limited range and low values of the covariances that their MAXCOV analyses generated. Consequently, they may not have scaled the ordinate of their graphs appropriately for the interpretation of the curves calculated from their data. A more extended ordinate makes their apparently peaked curve look flat and dimensional.

Later, Skilling, Quinsey, and Craig (2001) replicated Harris et al.'s (1994) results, using a similar methodology, but this time analyzing a sample of 1,111 boys. At first sight, their results appear to support the taxonic structure of psychopathy, but once again methodological problems plague this study. First, the authors unnecessarily dichotomized their items for MAXCOV, which weakens this procedure (Ruscio, 2000). Second, the authors used the goodness of fit index (GFI) to determine the nature of the latent structure. In the two studies that have examined the GFI systematically (Cleland, Rothschild, & Haslam, 2000; Haslam & Cleland, 2002), it has been shown to discriminate poorly between taxonic and dimensional structure. Moreover, because both taxonic and dimensional structures can yield GFIs well above .90, no universally applicable threshold has emerged even for data that GFI handles well. Examining the consistency of base rate estimates has intuitive appeal and is widely recommended and practiced, but nobody has ever actually established that taxonic structure does in fact yield more consistent estimates across

a realistic range of data parameters. A recent factorial Monte Carlo study (Ruscio, 2006) found that MAMBAC, MAXCOV, and MAXEIG analyses seldom yielded lower *SDs* for taxonic structure than for dimensional structure. More often than not, they were basically the same. Ambiguous MAMBAC and MAXCOV results may be produced by a positive skew of indicators or by low endorsement rates in the case of binary indicators. Rising MAMBAC curves and apparent (but low) peaks toward the right side of MAXCOV curves, highly consistent base rate estimates, and a high GFI are all as consistent with a latent dimension whose indicators are positively skewed (low endorsement), as they are with a small taxon.

Skilling, Harris, Rice, and Quinsey (2002) used the same participants as the 1994 study to investigate taxonic structure of APD, PCL-R, and the Child and Adolescent Taxon Scale (CATS), a derived instrument based on 8 items associated with the taxon in the Harris et al. (1994) paper. The authors performed MAMBAC and MAXCOV analyses and concluded that there was evidence supporting a taxon both for APD and for CATS. They failed to report the amplitude of the mean differences in MAMBAC and the covariance scores in MAXCOV. Other problems identified in the Harris et al. (1994) paper, such as possible rater file-review and graph comparison biases, absence of taxon base rate estimates, the potentially misleading admixed sample, and limited ordinate values also apply to the Skilling et al. (2002) study.

Recently, Vasey, Kotov, Frick, and Loney (2005) studied a sample of 386 children and adolescents to test the latent structure of psychopathy. The authors used two versions (Parent and Student) of the Antisocial Process Screening Device (APSD; Frick & Hare, 2001) to assess psychopathic characteristics. Along with evidence of a taxon for broad antisocial behavior, the authors claimed that they had found evidence for a psychopathy taxon. Specifically, using MAXEIG on the five subscales of youth self-report and parent APSD, the results produced

graphs consistent with a taxonic structure with an average base rate of .08. Their L-Mode analyses also suggested taxonicity with a base-rate of .04. Unfortunately, sample composition may easily explain the presence of the alleged taxon. In order to “increase the chances of detecting a psychopathy taxon (should one exist)” the authors added an extra 60 juvenile offenders to an already heterogeneous sample of 283 children and adolescents (boys and girls) recruited from middle schools, and 43 referred boys with severe emotional, behavioural, or learning problems. Vasey et al.’s (2005) two studies of children and adolescents were also limited by their lack of comparison curves for taxonic and dimensional data to help interpret their results. Some of their interpretations of curves as indicating taxonicity are questionable. Moreover, their strongest results came from data in which they combined separate community and clinical samples, increasing the potential for identifying pseudotaxa (Ruscio & Ruscio, 2004a). As the authors indicate, their best evidence identifies a taxon with a base rate that is too broad for psychopathy, and may be more accurately interpreted as an externalizing syndrome.

The four studies that profess to offer support for the hypothesis that the latent structure of psychopathy is taxonic do not provide convincing data to support this claim. Consequently, we conducted the present study in which Meehl’s (1995) taxometric method was applied to PCL-R ratings of 4,865 offenders sampled from multiple prison settings and assessed both by interviews and archival file review. In addition to a large representative sample, the putative psychopathy taxon was sufficiently large to afford powerful taxometric analyses, including the generation of simulated taxonic and dimensional data with which to compare the research data results. In the sample in the present study, 19% ($n = 927$) had a score of 30 or higher. The present sample also permitted the construction and analysis of several indicator sets with which to assess the latent structure of psychopathy and permitted detailed analyses of subsamples to examine and eliminate

potential sampling biases. Hare's (2003) four-factor solution to the PCL-R was used as the basis of the primary analyses, with composite scores for each factor serving as indicators. Additional analyses were performed using items within each of these four factors—as well as the affective-interpersonal features and impulsivity-antisocial behavior factors—as indicators to assess the taxonicity of each. Thus, the present study addressed the methodological shortcomings of prior studies and allowed a convincing test of the taxonicity of psychopathy and its components.

Method

Participants

The initial sample of participants consisted of 5408 male prison inmates incarcerated in North American institutions. Data were collected from 15 different samples across Canada (British Columbia, Ontario, and Québec) and the United States (Wisconsin and North Carolina).

Sample 1 comprised 322 male inmates of a federal medium-security institution in British Columbia, all serving sentences of two years or more. They had volunteered to take part in several research projects. Sample 2 was composed of 121 male inmates of a provincial prison in British Columbia, all serving sentences of less than two years. Sample 3 contained 369 male inmates of a minimum-security institution in Wisconsin (Kosson, Smith, & Newman, 1990). In sample 4, 106 male inmates of the Institute Phillippe-Pinel de Montréal in the province of Québec were evaluated with a French version of the PCL-R (Côté & Hodgins, 1991) just prior to conditional release from minimum, medium, or maximum security federal prisons. Sample 5 was composed of 87 inmates of a medium-security prison in Kingston, Ontario (Serin, 1991). Sample 6 was composed of 152 African-American male inmates (Kosson, Smith, & Newman, 1990). Sample 7 included 60 male inmates from a medium-security prison in North Carolina (Johnson, 1990). In Sample 8, 197 male inmates from a federal medium-security prison in British

Columbia were evaluated with the PCL-R. Sample 9 included 172 male inmates from a federal forensic psychiatric institution in British Columbia. Most were participants in a violent offender treatment program. Participants from Sample 10 were 1190 male inmates (526 White and 664 African-American) from a state medium-security prison in Wisconsin. In Sample 11, 320 male inmates (227 White, 73 Native; 220 sex offenders) from a federal medium security prison in British Columbia that houses violent offenders were interviewed (Porter, Fairweather, Drugge, Hervé, Birt, & Boer, 2000). Sample 12 was composed of 60 male inmates (rapists) in federal medium- and maximum-security institutions in Ontario (Brown & Forth, 1997). Sample 13 contained 185 male inmates of a medium-security institution in British Columbia. Sample 14 was composed of 427 male violent offenders in a federal medium-security institution in Ontario (Simourd & Hoge, 2000). Finally, sample 15 was composed of a representative sample of 1640 male offenders, admitted to a regional reception and assessment center of the Correctional Service of Canada.

The average age of the 2300 offenders for whom data were available was 31.1 ($SD = 9.3$; range from 17 to 81). In general, the reliability of the PCL-R ratings was quite acceptable. Cronbach's alpha was .85, and the intraclass correlation was .86 for a single rating and .92 for the average of two ratings. Intraclass Correlation Coefficients (ICC_1) for single ratings on Factor 1 and Factor 2 of the PCL-R were respectively .75 and .85 and for averaged ratings .85 and .92 (ICC_2). For the four facets (Hare, 2003), coefficients ranged from .67 to .84 for single ratings and from .80 to .91 for averaged ratings. To qualify as a PCL-R rater, individuals had to receive proper training in the use of the instrument. For a more detailed description of the sample composition and the rating procedures, see Hare (2003). Participants with missing information

on any one of the items ($n = 543$) were excluded, and analyses for the present study were performed on the 4865 participants with complete PCL-R protocols.

Measures

Psychopathy was assessed with the PCL-R (Hare, 1991). All participants were assessed using the semi-structured interview and file information. The PCL-R is a 20-item clinical rating scale that measures interpersonal, affective, and socially deviant features of psychopathy. The mean score on the PCL-R for the entire sample was 21.9 ($SD = 7.9$).

Taxometric Analyses

Two taxometric procedures were performed on each indicator set: MAMBAC (Mean Above Minus Below A Cut; Meehl & Yonce, 1994) and MAXEIG (MAXimum EIGenvalue; Waller & Meehl, 1998). A third procedure (L-Mode, for Latent Mode; Waller & Meehl, 1998) was performed only on the indicators in the primary analyses, which possessed sufficient variation to render a factor analysis meaningful. These three procedures are based on independent mathematical derivations and therefore can contribute nonredundant evidence of latent structure. Below, we note how MAMBAC and MAXEIG analyses were conducted, both of which involve a number of choice-points in their implementation. For detailed descriptions of the logic and mathematical underpinnings of all three procedures, see the primary publications cited above; on the available options for implementing each procedure and suggestions for making informed choices, see Ruscio and Ruscio (2004b, 2004c).

MAMBAC was performed using composite input indicators (i.e., selecting one of the k variables to serve as the output indicator and summing the remaining $k - 1$ variables to serve as the input indicator for each of k analyses). To accommodate the constrained indicator response scales, 50 equally-spaced cutting scores beginning and ending with at least 25 cases from each

end of the input indicator (larger values were used as necessary to stabilize the shapes near the ends of the curves) and 10 internal replications in the calculation of each curve (to reduce the obfuscating influence of cutting between equal-scoring cases) were used. MAMBAC curves were not smoothed, and full panels of curves for each indicator set were examined to determine whether an averaged curve adequately represented the overall trend. Averaged curves are presented here to conserve space. Panels are available upon request.

MAXEIG was performed using composite input indicators (in the same way, and for the same reason, as in MAMBAC), except for the primary series of analyses. In that instance, there was sufficient response variation to perform MAXEIG in the more traditional manner (i.e., selecting one of the k variables to serve as the input indicator and using the remaining $k - 1$ variables as output indicators for each of k analyses). Each analysis used 50 windows that overlapped 90% with adjacent subsamples and 10 internal replications. Once again, smoothing was not performed and full panels of curves were examined to ensure that the averaged curve fairly represented the overall pattern of results.

Analyses of Comparison Data

As a supplement to the inspection of taxometric results that can aid interpretation in ambiguous circumstances, bootstrap methods were used. This involves generating samples of taxonic and dimensional comparison data that reproduce the sample size, indicator distributions, and indicator correlations in a sample of research data and submitting each bootstrap sample to the same taxometric analyses as the research data. Comparison data were generated using an updated version of the algorithm outlined in Ruscio and Ruscio (2004a) and Ruscio, Ruscio, and Keane (2004) that implements two new features. First, comparison data are allowed to be multidimensional rather than constrained to unidimensionality (in the full sample for

dimensional comparison data, within groups for taxonic comparison data). Second, indicator distributions are generated using a standard bootstrap technique rather than being copied from the research data. The basic bootstrap approach of resampling, with replacement, treats a sample distribution as an unbiased estimate of the population distribution and draws new samples accordingly. Specifically, N scores for each indicator in a bootstrap sample of comparison data were drawn at random (with replacement) from the original score distribution. Programs including these features have been shown to reproduce indicator distributions and correlations with good precision and negligible bias (Ruscio, Ruscio, & Meron, 2005). Simulating taxonic comparison data requires a criterion variable that contains classification codes for each case. The PCL-R cutoff score of 30 that has typically been used for the diagnosis of psychopathy in research studies was for all analyses. This cutoff was originally generated to maximize the overall hit-rate of PCL-R for global judgments of psychopathy (cf. Hare, 2003). It has been used in hundreds of laboratory and applied research studies, and its utility has been supported by its generation and validation of many of the processing and neurological differences discussed in the introduction. There are also item response theory (IRT) data that suggest that the score of 30 indicates the same level of psychopathy across North American male offenders, English male offenders, female offenders, and male forensic psychiatric patients (Hare & Neumann, 2006). Consequently, it constitutes a reasonable cutoff for our taxonomic analyses. Because there might be some differences in offenders whose PCL-Rs were scored only from file reviews and those scored from a combination of file and interview assessments (Hare, 2003), we included in our study only protocols that were rated using both informational sources. We assigned the 927 cases with PCL-R total scores at or exceeding 30 to the putative taxon. We assigned all others to the

putative complement. We generated 10 samples of taxonic and 10 samples of dimensional comparison data for each analysis.

An examination of the results for taxonic and dimensional comparison data helps to ensure that apparently dimensional results do not stem from too small a sample size, inadequate representation of putative taxon members, insufficiently valid indicators, problematically high levels of nuisance covariance, or other unacceptable aspects of the data. The extent to which the results for the research data are better reproduced by those within the sampling distribution yielded by analyses of taxonic or dimensional comparison data sets supports the validity of the corresponding structural inference.

As an objective adjunct to the interpretation of curve shapes, we calculated a quantitative index. The fit of the averaged curve for the research data to the averaged curves in the sampling distributions of taxonic and dimensional comparison data sets was assessed using an approach introduced by Ruscio and Ruscio (2004b) and refined by Ruscio et al. (2005). First, fit is calculated for the comparison data generated to represent each structure:

$$Fit_{RMSR} = \sqrt{\frac{\sum (y_{res.data} - y_{sim.data})^2}{N}}, \quad (\text{Eq. 1})$$

where $y_{res.data}$ refers to a data point on the curve for the research data, $y_{sim.data}$ refers to the corresponding data point on the curve for comparison data, and N is the number of data points on each curve. Lower values of Fit_{RMSR} reflect better fit, with perfect fit represented by a value of 0. Equation 1 is calculated twice, once to assess the fit of the research curves to those for taxonic comparison data ($Fit_{RMSR-tax}$) and once to assess the fit for dimensional comparison data ($Fit_{RMSR-dim}$). Then, these two values are integrated into a single comparison curve fit index (CCFI):

$$CCFI = \frac{Fit_{RMSR-dim}}{Fit_{RMSR-dim} + Fit_{RMSR-tax}}, \quad (\text{Eq. 2})$$

CCFI values can range from 0 to 1, with lower values suggesting better fit for dimensional structure and higher values suggesting better fit for taxonic structure. The index is symmetrical about .50 in that this middle value represents equivalent fit for both structures. It is important to note that the CCFI indexes the relative fit of taxonic and dimensional structural models, not the absolute goodness of fit of either model. In a preliminary test of this type of curve-fit index (Ruscio, 2004), latent structure was correctly classified with high precision in analyses of the 700 Meehl and Yonce (1994) samples of taxonic and dimensional data. In a Monte Carlo study that include a much broader range of data conditions (Ruscio et al., 2005), this index significantly outperformed several of the most popular taxometric consistency tests.

Results

In the first series of analyses, composite indicators were formed in accordance with Hare's (2003) four-factor model; items assigned to each factor (referred to here as facets) were summed to yield one indicator apiece. These analyses are primary in the sense that the structure of psychopathy was represented by all four facets: Interpersonal, Affective, Lifestyle, Antisocial. The results for MAMBAC, MAXEIG, and L-Mode analyses of the four summative indicators appear in Figure 1. These analyses yielded quite clear and consistent results. Curve shapes were highly consistent with what one would expect for dimensional structure, as well as the curves observed for simulated dimensional comparison data (which is also reflected in CCFI values that strongly favor a dimensional interpretation; see Table 1 for all CCFI results). Although there was a modest level of consistency among the MAMBAC base rate estimates, as well as among the MAXEIG estimates, this was not true of the L-Mode estimates (see Table 2 for summaries of all taxon base rate estimates). In addition to the notable discrepancies across procedures' estimates, the results were much better reproduced by the dimensional than the taxonic comparison data.

Thus, the totality of evidence suggests that these four theoretically-derived composite indicators of psychopathy do not represent a taxonic construct. Rather, individual differences appear more consistent with dimensional structure.

The second series of taxometric analyses was performed to test the latent structure of each of the four theoretical facets of psychopathy. The results for MAMBAC and MAXEIG analyses of indicator sets representing each facet (Affective, 4 indicators; Antisocial, 5 indicators; Interpersonal, 4 indicators; and Lifestyle, 5 indicators) appear in Figure 2. In analyses of the research data, no taxonic peaks emerged, and CCFI values supported—in 7 out of 8 analyses—an inference of dimensional structure for each facet. Taxon base rate estimates varied substantially within analyses, and they were generally better reproduced by the dimensional than the taxonic comparison data. Thus, although the results are not as prototypically dimensional in appearance as in the first series of analyses, no evidence in support of a taxon was obtained for any of the four theoretical facets.

The final series of taxometric analyses examined the latent structure of PCL-R Factors 1 and 2 using separate sets of indicators to represent each. The results for MAMBAC and MAXEIG analyses of indicator sets representing PCL-R Factors 1 (8 indicators) and 2 (10 indicators) appear in Figure 3. None of the curves for the research data contain the peaks that would be expected of taxonic data, an interpretation supported by CCFI values in 3 out of 4 analyses (the exception was an ambiguous value supportive of neither structure). Moreover, taxon base rate estimates were inconsistent within and across procedures. Results for comparison data suggest that either latent structure could have given rise to such estimates. Hence they are not particularly informative in these analyses. To the extent that the results for these two factors hint at latent structure, they suggest a dimensional interpretation. In any event, there is no

evidence that either the Affective/Interpersonal (Factor 1) or the Lifestyle/Antisocialty (Factor 2) features of the PCL-R factor are distributed as a taxon.

Follow-up analyses were performed for each indicator set within subsamples consisting of (a) inmates in maximum security prisons, (b) inmates in medium security prisons, (c) inmates in minimum security prisons, (d) black inmates, (e) white inmates, and (f) and inmates from a representative sample of subjects incarcerated in Canadian federal institutions. These analyses yielded far too many curves to present here, but all CCFI values are provided in Table 1. For each indicator set, results across subsamples were consistent with those in the full sample in supporting an inference of dimensional structure.

Discussion

Summary of Results

Overall, the observed results support a dimensional structure for psychopathy and its components as measured by the PCL-R. Whether considered separately or in combination, none of the analyses provided results consistent with a taxonic structure. The primary analyses—those performed using four theoretically-based composite indicators—yielded unambiguously dimensional results, and the follow-up analyses produced results that did not follow the patterns observed for prototypical taxonic or dimensional data. Nonetheless, we believe that the totality of evidence warrants a dimensional interpretation for several reasons. First, none of the MAMBAC or MAXEIG curves generated peaks supportive of a taxonic structure. Second, a curve-fit index based on bootstrapped sampling distributions of results for taxonic and dimensional comparison data consistently suggested dimensional structure. Third, taxon base rate estimates were inconsistent across analyses and, more often than not, the *M*s and *SD*s of

these estimates were better reproduced by the dimensional than the taxonic comparison data.

Fourth, regardless of how the sample was divided, these same results were found.

Integration of Current Results with Previous Studies

These results are consistent with the results found by Guay and Knight (2003) using the MASA and by Marcus et al. (2004) using the PPI, but differ from the four studies that identified taxonic distributions. It is likely that the methodological problems in those four studies, delineated in the introduction and addressed in the present study, are responsible for the differences in the results. The present study countered the problems of these prior studies by accessing a larger and more representative sample and analyzing in detail subsamples to assure that particular selection criteria did not bias results. Both interview and archival file review were employed in arriving at PCL-R scores, and only those offenders for whom ratings on all items were possible were included, guaranteeing a more complete and detailed coverage of both Factor 1 and Factor 2 content and guarding against the biasing tendency for raters to use related items to arrive at judgments for items with insufficient information. Finally, more sophisticated analytic procedures were introduced, including the comparison of simulated taxonic and dimensional comparisons generated from the same sample and quantitative analyses of curve fits to reduce some of arbitrariness of the interpretation of the results.

Interpretations of the Dimensional Results

The clear and strong indications of the dimensionality of psychopathy can be interpreted in several ways. These range from the acceptance of psychopathy as dimensional and possibly an extreme on one or more normative personality traits to questioning the PCL-R as the appropriate assessment to uncover a taxon, to considerations about the homogeneity of psychopathy and the

potential multiplicity of its underlying core processes. We will consider each of these alternative perspectives in turn.

Psychopathy as a dimension. Consistent with the results of these taxometric analyses we might conclude that both the latent structure of psychopathy and its core factors may best be interpreted as dimensionally distributed. A shortcoming of taxometric analyses is its failure to parameterize any particular alternative dimensional model (Krueger, 2006). Consequently, the disconfirmation of a taxonic model does not suggest or corroborate any particular alternative model, but rather is simply consistent with research that supports the hypothesis that personality disorders in general may be best conceptualized as distinct configurations of extreme scores on personality traits, affective and cognitive competences, or neurobiological processes that exist on a continuum with normal functioning (e.g., Widiger, 1993; Widiger & Costa, 1994). Using the perspective of the five-factor model (FFM; McCrae & Costa, 1990), Lynam (2002) has recently presented data that support the application of a dimensional conceptualization to psychopathy. He argued both from expert-generated FFM psychopathy prototypes and correlations of the FFM with measures of psychopathy that the psychopath could be described as low on all facets of Agreeableness, and low in the dutifulness, deliberation, and self-discipline facets of Conscientiousness. Mixed results emerged for Neuroticism and Extraversion. For the psychopath whereas the Neuroticism facets of anxiety, self-consciousness, and perhaps vulnerability and depression were low, the facets of impulsivity and angry hostility were high. The Extraversion facet of excitement seeking was high, but the facets of warmth and perhaps positive emotions were low. Lynam (2002) concluded that psychopathy could best be conceptualized by extreme scores on a collection of FFM personality traits. More generally, Bishopp & Hare (2005) have reported that a multidimensional scaling analysis of the data set used in the present study

provided support for a multidimensional structure within the PCL-R, corroborating the hypothesis that psychopathy can be viewed as an extreme variant of multiple dimensions of personality. Psychopathy might also be conceptualized within the framework of an “externalizing spectrum” of personality and psychopathology (Krueger, 2006).

Conceiving of psychopathy as a dimension carries several implications about optimal strategies for studying the disorder. It argues for moving away from extreme group designs that attempt to distinguish psychopaths from nonpsychopaths or from trichotomizing PCL-R scores and assuming that discrete groups have been formed for comparison purposes (Lilienfeld, 1994), and toward dimensional designs, such as the quantitative, latent trait model-based approach proposed by Krueger, Markon, Patrick, and Iacono (2005) in explicating the comorbidity among externalizing disorders. The dimensionalization of psychopathy is also consistent with the recent increase in research on subclinical manifestations of psychopathy (Hall & Benning, 2006) and suggests the importance of such research for unravelling etiological factors of the components of psychopathy. In this regard it is interesting to note that the factors of psychopathy have recently been identified as critical in predicting sexual coercion against women (Knight & Guay, 2006). The predictive potency of these factors is similar in criminal and non-criminal, juvenile and adult samples (Knight & Sims-Knight, 2003; 2004) suggesting a gradual rather than a step function in the contribution of psychopathy to rape.

Certainly, the strong evidence for the dimensional latent structure of psychopathy should affect how this construct is conceptualized and used in criminal justice proceedings. The PCL-R is the most commonly cited measure in such proceedings (Hare, 1996; Edens & Petrila, 2006) and is a part of some actuarials that have been constructed to predict both violence (e.g., Violence Risk Appraisal Guide; Harris, Rice, & Quinsey, 1993) and sexual violence (e.g., Sex

Offender Risk Appraisal Guide; Quinsey et al., 1998). In civil commitment procedures for sexual predators it is sometimes a critical issue to determine whether a defendant is or is not “a psychopath” (Edens & Petrila, 2006). The present research suggests that criminal justice language and conceptualization should be modified and should stop talking about individuals as being “psychopaths.” Rather, lawyers should refer to defendants as being “high on measures of psychopathy.” Although this appears to be a subtle difference, it may have important consequences in reducing juries’ perceptions of particular defendants as different in kind. The suggestion of Edens and Petrila (2006) to have confidence intervals as well as discrete scores reported in court cases is also apropos. The data supporting the dimensionalization of psychopathy also suggest that actuarial measures used in the criminal justice system might profit from using the full PCL-R score rather than reifying an arbitrary dichotomization or weighting scores to give more leverage to a purported taxon.

PCL-R as a questionable intervening measure. The specificity both of the PCL-R and also self-report measures of psychopathy like the MASA and PPI may not be sufficient to identify genotypic and biological causation. It is possible that such assessment instruments, which predominantly measure interpersonal behavior in a social environment, may be too distant phenotypically from the biological substrate to covary with a hypothetical taxon. Other phenotypic indicators should be explored. One approach might be to use as dependent variables scores on the cognitive (e.g., Kosson, 1996; 1998; Newman, 1987; Newman et al., 1987; Schmauk, 1970) and affective (e.g., Levenston, Patrick, Bradley, & Lang, 2000; Patrick, Bradley, & Lang, 1993) measures that have been found to covary with psychopathy and might better tap processes more closely related to underlying mechanisms. Less molar, less “psychological” or “social” kinds of indicators are likely to be connected to any putative

underlying disposition by shorter causal chains and hence may involve fewer attenuating factors. Such measures may have a higher likelihood of being taxometrically strong indicators of psychopathy, if indeed it constitutes a taxon.

Heterogeneity in psychopathy and the obscuring role of subtypes. The typological purity of psychopathy has often been challenged. It has most commonly been proposed that two distinct subtypes of psychopaths can be identified, often referred to as primary and secondary psychopathy (Skeem, Poythress, Edens, Lilienfeld, & Cale, 2003). Frick, Lilienfeld, Ellis, Loney, and Silverthorn (1999) have argued that “cooperative suppressor” effects of two correlated psychopathic types might mask important differences between the types. It might also be argued that the presence of these correlated subtypes might mask taxonic differences as well.

Using Fraley’s (1998) model-based clustering, which in contrast to other clustering techniques provides a fit index that allows one to assess the best fitting model, Hicks, Markon, Patrick, Krueger, & Newman (2004) have provided the most compelling demonstration of these subtypes. They cluster analyzed a group of high PCL-R psychopaths, using the scales from the Multidimensional Personality Questionnaire (MPQ-BF; Patrick, Curtin, & Tellegen, 2002) as dependent measures. The best model yielded two clear clusters. The first cluster, comparable to the primary psychopath and labeled the stable psychopath, was low in Stress Reaction, Social Closeness, and Harm Avoidance, and high in Agentic positive emotionality. The authors interpreted this type as approximating the classical psychopath, who is immune to stress, socially dominant, but unattached to others, and prone to take risks. The second subtype, called the aggressive psychopath, scored high on Negative Emotionality and low on Constraint and was characterized by high aggression. This type appears to capture undercontrolled, externalizing

individuals (e.g., Krueger et al., 2002), whom the authors compared to Moffitt's (1993) life-course persistent offenders.

Even though these markedly different personality types apparently emphasize specific characteristics of the PCL-R superordinate factors, nonetheless these types are not congruent with these factors, and their presence might mask the manifestation of underlying taxa. One strategy to resolve this issue would be to use measures independent of the PCL-R to remove one or the other of these subtypes and conduct the taxometrics on the remaining offenders. For instance, one could follow Newman's strategy (e.g., Newman & Schmitt, 1998; Newman et al., 1997) and use the Welsh Anxiety Scale (Welsh, 1956) to remove alternatively high- and low-anxious psychopaths from the analyses. The results of Hicks et al. (2004) could also be used to identify MPQ-BF profiles for the two subtypes, and each subtype could be removed in alternate analyses leaving the other to determine whether a taxon solution emerges in the absence of either.

Problems of mapping from the phenotypic to underlying causes. A final possible interpretation is illustrated in the neurobehavioral model of personality traits proposed by Depue and Lenzenweger (2001). They conceive personality disorders as functions of the variations or interactions of the most extreme values that contribute to high-order traits: Constraint, Affiliation, Positive-Negative Emotionality (PEM-NEM), and Fear. The authors propose that higher-order heterogeneous phenotypes result from the combination of a heterogeneous set of lower-order traits that have different sources of genetic variation. The task of identifying neurobiological foundations is rendered particularly difficult, considering that some higher-order traits are typically associated with two or more behavioral systems and neurobehavioral networks

(Depue & Lenzenweger, 2001). Consequently, it would be particularly difficult to observe taxonic results.

As Gottesman (1997) illustrates in the domain of intelligence assessment, phenotypical continuity can represent the true manifestation of distinct underlying genotypic distributions. Genetic and biological causation and phenotypic discontinuity are not synonymous. Depue and Lenzenweger (2001) suggest that at least four different neurobehavioral systems may be necessary to account for impulsivity—(a) positive incentive motivation; (b) fear; (c) aggression; and (d) low levels of a nonaffective form of impulsivity (which results in disinhibition of the above neurobehavioral systems). For impulsivity to emerge at least four to five independent neurobehavioral systems may have to be elicited (not to mention an increased complexity when possible interaction with other higher-order traits is considered). Ultimately, the determination of the number of core processes that are necessary and sufficient to explain and predict psychopathy will depend on the isolation of the basic processes involved (Depue & Lenzenweger, 2001; Tellegen & Waller, in press). Specific etiologies, including particular genetic origins, prenatal, perinatal, and postnatal biological determinants, and specific life experiences impacting on causal neurobiological influences within personality structure will determine the relevant processes contributing to psychopathy. As in the hunt for the “quantitative trait loci” of intelligence (Gottesman, 1997), such complexity does not easily yield its underlying causal mechanisms to empirical scrutiny, and investment in a long-term search is necessary. The advantage of being able to identify a taxon in the phenotypical distributions of PCL-R or its subcomponents would have been the location of a beacon guiding us to a specific etiology (Meehl, 1977). The absence of such a beacon may mean that no phenotypic measures, even cognitive or affective assessments that may be closer to the underlying causal mechanisms, will

provide such guidance and longitudinal research strategies aimed at identifying and tracking multiple underlying processes and environmental contingencies will be necessary.

Conclusions

The analyses in the present paper clearly indicate that the disorder defined by high scores on the PCL-R and the correlated factors that the PCL-R comprises are distributed dimensionally. Although such results strongly disconfirm the hypothesis that the PCL-R covaries with a latent taxon, questions about the specificity of the measure in covarying with underlying process, the possibility of covarying, but etiologically distinct subtypes of psychopathy, and the possibility that the manifest behavior of psychopathy is the interaction of multiple independent neurobiological systems leave open the possibility that other measures might yet uncover a taxon. The support for a dimensional model of psychopathy buttresses research strategies that are examining the correlates of psychopathy self-report scales in both criminal and non-criminal samples (Benning, Patrick, Salekin, & Leistico, 2005; Brook, Kosson, Walsh, & Robins, 2005; Knight & Sims-Knight, 2003), encourages structural equation modeling approaches to conceptualizing and testing etiological hypotheses about psychopathy (Krueger et al., 2002), and argues for the creation of scales that are equally reliable and discriminating across the full spectrum of the scale. Although we should not yet forego the more traditional group comparison approaches, such data should be analyzed and interpreted with dimensional precautions in mind. We have not resolved the issue of taxonicity sufficiently to leave any strategy behind in the attempts to unravel the Gordian knot of psychopathy.

References

- Alpert, M., Shaw, R. J., Pouget E. R., & Lim K. O. (2002). A comparison of clinical ratings with vocal acoustic measures of flat affect and alogia. *Journal of Psychiatric Research* 36, 347–353.
- American Psychiatric Association. (1994). *Diagnostic and Statistical Manual of Mental Disorders*, 4th edition, Washington, D.C.
- Beauchaine, T. P., & Waters, E. (2003). Pseudotaxonomicity in MAMBAC and MAXCOV analyses of rating scale data: Turning continua into classes by manipulating observer's expectations. *Psychological Methods*, 8, 3-15.
- Bennett, A. J., Lesch, K. P., Heils, A., Long, J.C., Lorenz, J.G., Shoaf, S.E., Champoux, M., Suomi, S.J., Linnoila, M.V., & Higley, J. D. (2002). Early experience and serotonin transporter gene variation interact to influence primate CNS function, *Molecular Psychiatry*, 7, 118–122
- Benning, S. D., Patrick, C. J., Salekin, R. T., & Leistico, A. R. (2005). Convergent and discriminant validity of psychopathy factors assessed via self-report: A comparison of three instruments. *Assessment*, 12, 270-289.
- Benning, S. D., Patrick, C. J., Blonigen, D. M., Hicks, B. M., & Iacono, W. G. (2005). Estimating facets of psychopathy from normal personality traits: A step toward community-epidemiological investigations. *Assessment*, 12, 3-18.
- Bishopp, D., & Hare, R. D. (2005). *A multidimensional scaling analysis of the Hare PCL-R*. Manuscript submitted for publication.
- Blair, R. J. R. (1995). A cognitive developmental approach to morality: Investigating the psychopath. *Cognition*, 57, 1-29.

- Blair, R. J. R. (2001). Neuro-cognitive models of aggression, the antisocial personality disorders and psychopathy. *Journal of Neurology, Neurosurgery and Psychiatry*, *71*, 727 -731.
- Blair, R. J. R. (2004). The roles of orbital frontal cortex in the modulation of antisocial behavior. *Brain and Cognition*, *55*, 198-208.
- Blair, R. J. R., Colledge, E., Murray, L., & Mitchell, D. G. V. (2001). A selective impairment in the processing of sad and fearful expressions in children with psychopathic tendencies. *Journal of Abnormal Child Psychology*, *29*, 491-498.
- Blair, R. J. R., Mitchell, D. G. V., Richell, R. A., Kelly, S., Leonard, A., Newman, C. & Scott, S. K. (2002). Turning a deaf ear to fear: Impaired recognition of vocal affect in psychopathic individuals. *Journal of Abnormal Psychology*, *111*, 682-686.
- Blair, R. J. R., Mitchell, D. G. V., & Blair, K. S. (2005). *The psychopath: Emotion and the brain*. Oxford: Blackwell.
- Blonigen, D. M., Carlson, S. R., Krueger, R. F., & Patrick, C. J. (2003). A twin study of self-reported psychopathic personality traits. *Personality and Individual Differences*, *35*, 179-197.
- Blonigen, D. M., Hicks, B. M., Kreuger, R. F., Patrick, C. J. & Iacono, W. G. (2005). Psychopathic personality traits : heritability and genetic overlap with internalizing and externalizing psychopathology. *Psychological Medicine* , *35*, 1-12.
- Bolt, D., Hare, R. D., Vitale, J., & Newman, J. P. (2004). A multigroup item response theory analysis of the Hare Psychopathy Checklist—Revised. *Psychological Assessment*, *16*, 155-168.
- Brook, M., Kosson, D. S., Walsh, Z., & Robins, R. W. (2005, July). *Psychopathic personality as measured by the Self-Report Psychopathy Scale-II (SRP-II): External correlates,*

- antisocial deviance, and substance use*. Poster presented at the Annual Convention of the Society for the Scientific Study of Psychopathology. Vancouver, Canada.
- Cadore, R. J., Yates, W. R., Troughton, E., Woodworth, G., & Stewart, M. A. (1995). Adoption study demonstrating two genetic pathways to drug abuse. *Archives of General Psychiatry*, *42*, 161-167.
- Cleckley, H. (1976). *The mask of sanity* (5th ed). St Louis: C.V. Mosby.
- Cleland, C., Rothschild, L., & Haslam, N. (2000). Detecting latent taxa: Monte Carlo comparison of taxometric, mixture and clustering methods. *Psychological Reports*, *87*, 37-47.
- Cooke, D. J., & Michie, C. (2001). Refining the construct of psychopathy: Towards a hierarchical model. *Psychological Assessment*, *13*, 171-188.
- Depue, R. A., (1996). A neurobiological framework for the structure of personality and emotion: Implications for personality disorders. In J. F. Clarkin & M. F. Lenzenweger (Eds.), *Major theories of personality disorder* (pp. 347–390). New York: Guilford Press.
- Depue, R. & Lenzenweger, M. (2001). Neurobiology of personality disorders. In J. Livesley (Ed.), *Handbook of Personality Disorders* (pp. 136-176). New York: Guilford Press.
- Douglas, K., Vincent, G. M., & Edens, J. (2006). Risk for Criminal Recidivism: The role of psychopathy. In C. Patrick (Ed.), *Handbook of Psychopathy* (pp. 533-554). New York: Guilford Press.
- Edelbrock, C., Rende, R., Plomin, R., & Thompson, L. A. (1995). A twin study of competence and problems behavior in childhood and early adolescence. *Journal of Child Psychology and Psychiatry*, *36*, 775-785.

- Edens & Petrila, (2006). Legal and Ethical Issues in the Assessment and Treatment of Psychopathy. In C. Patrick (Ed.), *Handbook of Psychopathy* (pp. 573-588). New York: Guilford Press.
- Fowles, D. C. & Dindo, L. (2006). A dual-deficit model of psychopathy. In C. Patrick (Ed.), *Handbook of Psychopathy* (pp. 14-34). New York: Guilford Press.
- Fraley, C. (1998). Algorithms for model-based Gaussian hierarchical clustering. *Journal on Scientific Computing*, 20, 270-281.
- Frick, P. J., & Hare, R. D. (2001). *The antisocial process screening device*. Toronto: Multi-Health Systems.
- Frick, P. J., Lilienfeld, S. O., Ellis, M. L, Loney, B. R., & Silverthorn, P. (1999). The association between anxiety and psychopathy dimensions in children. *Journal of Abnormal Child Psychology*, 27, 381-390.
- Ge, X., Conger, R. D., Cadoret, R. D., Neiderhiser, J., Troughton, E., Stewart, E., & Yates, W. (1996). The developmental interface between nature and nurture: A mutual influence model of adolescent antisocial behavior and parenting behaviors. *Developmental Psychology*, 32, 574-589.
- Gottesman, I. I. (1997). Twins - en route to QTLs for cognition. *Science*, 276, 1522-1523.
- Gottesman, I. I., & Goldsmith, H. H. (1994). Developmental psychopathology of antisocial behavior: Inserting genes into its ontogenesis and epigenesis. In C. A. Nelson (Ed.), *Threats to optimal development, integrating biological, psychological, and social risk factors* (pp. 69-104). Hillsdale, NJ: Lawrence Erlbaum.

- Grove, W. M., Eckert, E. D., Heston, L., Bouchard, T. J., Jr., Segal, N., Lykken, D. T. (1990). Heritability of Substance Abuse and Antisocial Behavior: A Study of Monozygotic Twins Reared Apart. *Biological Psychiatry*, 27, 1293-1304.
- Guay, J. P., & Knight, R. A. (2003, July). *Assessing the Underlying Structure of Psychopathy Factors using Taxometrics*. Poster presented at the Developmental and Neuroscience Perspectives on Psychopathy conference, Madison, Wisconsin.
- Hall, J. R. & Benning, S. D. (2006). The "Successful" Psychopath: Adaptive and Subclinical Manifestations of Psychopathy in the General Population. In C. Patrick (Ed.), *Handbook of Psychopathy* (pp. 459-478). New York: Guilford Press.
- Hare, R.D. (1980). A research scale for the assessment of psychopathy in criminal populations. *Personality and Individual Differences*, 1, 111–119.
- Hare, R. D. (1991). *The Psychopathy Checklist-Revised manual*. Toronto, Canada: Multi-Health Systems.
- Hare, R. (1996). Psychopathy: A clinical construct whose time has come. *Criminal Justice and Behavior*, 23, 25-54.
- Hare, R. D. (2001). Psychopaths and their nature: Some implications for understanding human predatory violence. In A. Raine & J. Sanmartin (Eds.), *Violence and Psychopathy* (pp. 5-34). Dordrecht, The Netherlands: Kluwer Academic Publishing.
- Hare, R. D. (2003). *The Psychopathy Checklist-Revised technical manual* (2nd ed.). Toronto, Canada: Multi-Health Systems.
- Hare, R. D., Harpur, T. J., Hakstian, A. R., Forth, A. E., Hart, S. D., & Newman, J. P. (1990). The revised Psychopathy Checklist: Reliability and factor structure. *Psychological Assessment* 2, 338-341.

- Hare, R. D., & Neumann, C. N. (2006). The PCL-R Assessment of Psychopathy: Development, Structural Properties, and New Directions. In C. Patrick (Ed.), *Handbook of Psychopathy* (pp. 58-88). New York: Guilford Press.
- Harpur, T. J., Hakstian, A. R., & Hare, R. D. (1988). Factor structure of the Psychopathy Checklist. *Journal of Consulting and Clinical Psychology, 56*, 741-747.
- Harpur, T. J., Hare, R. D., & Hakstian, A. R. (1989). Two-factor conceptualization of psychopathy: Construct validity and assessment implications. *Psychological Assessment, 1*, 6-17.
- Harris, G. T., Rice, M. E., & Quinsey, V. L. (1994). Psychopathy as a taxon: Evidence that psychopaths are a discrete class. *Journal of Consulting and Clinical Psychology, 62*, 387-397.
- Hart, S. D., Kropp, P. R., & Hare, R. D. (1988). Performance of male psychopaths following conditional release from prison. *Journal of Consulting and Clinical Psychology, 56*, 227-232.
- Hart, S. D., & Hare, R. D., (1989). Discriminant Validity of the Psychopathy Checklist in a forensic psychiatric population. *Journal of Consulting and Clinical Psychology, 1*, 211-218.
- Hart, S. D., & Hare, R. D. (1997). Psychopathy: Assessment and association with criminal conduct. In D. Stoff, J. Breiling, & J. Maser (Eds.), *Handbook of Antisocial Behavior* (pp. 22-35). New York, New York: Wiley & Sons.
- Haslam, N., & Cleland, C. (2002). Taxometric analysis of fuzzy categories: A Monte Carlo study. *Psychological Reports, 90*, 401-404.

- Hervé, H. F., Hayes, P. J., & Hare, R. D. (2003). Psychopathy and sensitivity to the emotional polarity of metaphorical statements. *Personality and Individual Differences, 35*, 1497-1507.
- Hicks, B. M., Markon, K. E., Patrick, C. J., Krueger, R. F., & Newman, J. P. (2004). Identifying psychopathy subtypes based on personality structure. *Psychological Assessment, 16*, 276-288.
- Hill, C., Neumann, C., S., & Rogers, R. (2004). Confirmatory factor analysis of the Psychopathy Checklist: Screening Version (PCL: SV) in offenders with Axis I Disorders. *Psychological Assessment, 16*, 90-95.
- Johnson, T. D. (1990). *Partial helplessness conditioning as a possible etiological factor in psychopathy*. Unpublished doctoral dissertation. University of Alabama, Birmingham, AL.
- Kiehl, K. A., Bates, A. T., Laurens, K. R., Hare, R. D., & Liddle, P. F. (in press). Brain potentials implicate temporal lobe abnormalities in criminal psychopaths. *Journal of Abnormal Psychology*.
- Kiehl, K. A., Hare, R. D., Liddle, P. F., & McDonald, J. (1999). Reduced P300 responses in criminal psychopaths during a visual oddball task. *Biological Psychiatry, 45*, 1498-1507.
- Kiehl, K. A., Smith, A. M., Hare, R. D., Mendrek, A., Forster, B. B., Brink, J., & Liddle, P. F. (2001). Limbic abnormalities in affective processing by criminal psychopaths as revealed by functional magnetic resonance imaging. *Biological Psychiatry, 50*, 677-684.
- Kiehl, K. A., Smith, A. M., Mendrek, A., Forster, B. B., Hare, R. D., & Liddle, P. F. (2004). Temporal lobe abnormalities in semantic processing by criminal psychopaths as revealed

- by functional magnetic resonance imaging. *Psychiatry Research: Neuroimaging*, 130, 27-42.
- Knight, R.A. (1995). *A unified theory of sexual aggression*. Paper presented at the 14th Annual Meeting of the Association for the Treatment of Sexual Abusers, New Orleans, LA.
- Knight, R.A., & Cerce, D.D. (1999). Validation and revision of the Multidimensional Assessment of Sex and Aggression. *Psychologica Belgica*, 39, 135-161.
- Knight, R.A. & Guay, J.P. (2006). The Role of Psychopathy in Sexual Coercion against Women. In C. Patrick (Ed.), *Handbook of Psychopathy* (pp. 512-532). New York: Guilford Press.
- Knight, R. A., & Sims-Knight, J. E. (2003). Developmental antecedents of sexual coercion against women: Testing of alternative hypotheses with structural equation modeling, In R. A. Prentky, E. Janus, & M. Seto (Eds.), *Sexual coercion: Understanding and management* (pp.72-85). New York: New York Academy of Sciences.
- Knight, R. A., & Sims-Knight, J. E. (2004). Testing an etiological model for juvenile sexual offending against women. *Journal of Child Sexual Abuse*, 13, 33-55.
- Kosson, D. S. (1996). Psychopathy and dual -task performance under focusing conditions. *Journal of Abnormal Psychology*, 105, 391-400.
- Kosson, D. S. (1998). Divided visual attention in psychopathic and nonpsychopathic offenders. *Personality and Individual Difference*, 24, 373-391.
- Kosson, D. S., Smith, S. S., & Newman, J. P. (1990). Evaluation of the construct validity of psychopathy in Black and White male inmates: Three preliminary studies. *Journal of Abnormal Psychology*, 99, 250-259.
- Krueger, R. F. (1999). The structure of common mental disorders. *Archives of General Psychiatry*, 56, 921-926.

- Krueger, R. F. (2000). Phenotypic, genetic, and nonshared environmental parallels in the structure of personality: A view from the Multidimensional Personality Questionnaire. *Journal of Personality and Social Psychology*, *79*, 1057-1067.
- Krueger, R.F. (2006) Perspectives on the Conceptualization of Psychopathy: Toward an Integration. In C. Patrick (Ed.), *Handbook of Psychopathy* (pp. 193-202). New York: Guilford Press.
- Krueger, R. F., Hicks, B. M., Patrick, C. J., Carlson, S. R., Iacono, W. G., & McGue, M. (2002). Etiologic connections among substance dependence, antisocial behavior, and personality: Modeling the externalizing spectrum. *Journal of Abnormal Psychology*, *111*, 411-424.
- Krueger, R. F., Markon, K. E., Patrick, C. J., & Iacono, W. G. (2005). Externalizing psychopathology in adulthood: A dimensional-spectrum conceptualization and its implications for DSM-V. *Journal of Abnormal Psychology*, *114*, 537-550.
- Larsson, H., Andershed, H., & Lichtenstein, P. (in press). A genetic factor explains most of the variation in the psychopathic personality. *Journal of Abnormal Psychology*.
- Lesch, K. P., Merschdorf, U. (2000). Impulsivity, aggression, and serotonin: a molecular psychobiological perspective. *Behavioral Sciences and the Law*, *18*, 581-604.
- Levenston, G. K., Patrick, C. J., Bradley, M. M., & Lang, P. J. (2000). The psychopath as observer: Emotion and attention in picture processing. *Journal of Abnormal Psychology*, *109*, 373-385.
- Lilienfeld S.O. (1994). Conceptual problems in the assessment of psychopathy. *Clinical Psychology Review*, *14*, 17-38.

- Lilienfeld, S. O., & Andrews, B. P. (1996). Development and preliminary validation of a self-report measure of psychopathic personality traits in noncriminal populations. *Journal of Personality Assessment, 66*, 488-524.
- Livesley, W. J. (1998). Suggestions for a framework for an empirically based classification of personality disorder. *Canadian Journal of Psychiatry, 43*, 137-147.
- Lynam, D. R. (2002). Psychopathy from the perspective of the Five Factor Model. In P. T. Costa, Jr., & T. A. Widiger (Eds.), *Personality disorders from the perspective of the Five Factor Model of personality* (2nd ed., 325-350). Washington, DC: American Psychological Association.
- Lyons, M. J., True, W. R., Eisen, S. A., Goldberg, J., Meyers, J. M., Faraone, L. J., Eaves, L. J., & Tsuang, M. T. (1995). Differential heritability of adult and juvenile antisocial traits. *Archives of General Psychiatry, 52*, 906-915.
- Marcus, D. K., John, S. L. & Edens, J. F. (2004). A Taxometric Analysis of Psychopathic Personality. *Journal of Abnormal Psychology, 113*, 626-635.
- Mason, D. A. & Frick, P. J. (1994). The heritability of antisocial behavior: A meta-analysis of twin and adoption studies. *Journal of Psychopathology and Behavioral Assessment, 16*, 301-323.
- McCrae, R. R., & Costa, P. T. (1990). *Personality in Adulthood*. New York: Guilford Press.
- Mednick, S. A., Gabrielli, W. F., Jr. & Hutchings, B. (1984). Genetic influences in criminal convictions: Evidence from an adoption cohort. *Science, 224*, 891-894.
- Meehl, P. E. (1973). MAXCOV-HITMAX: A taxonomic search method for loose genetic syndromes. In P.E. Meehl (Ed.), *Psychodiagnosis: Selected papers*, (pp. 200-224), Minneapolis: University of Minnesota Press.

- Meehl, P. E. (1977). Specific etiology and other forms of strong influence: Some quantitative meanings. *Journal of Medicine and Philosophy*, 2, 33-53.
- Meehl, P. E. (1992). Factors and taxa, traits and types, differences of degree and differences in kind. *Journal of Personality*, 60, 117-174.
- Meehl, P. E. (1995). Bootstraps taxometrics: Solving the classification problem in psychopathology. *American Psychologist*, 50, 266-275.
- Meehl, P. E. & Yonce, L. J. (1994). Taxometric analysis: I. Detecting taxonicity with two quantitative indicators using means above and below a sliding cut (MAMBAC procedure). *Psychological Reports*, 74, 1059-1274.
- Miller, J. D., Lynam, D. R., Widiger, T. A., & Leukefeld, C. (2001). Personality disorders as extreme variants of common personality dimensions. Can the Five-factor model of personality adequately represent psychopathy? *Journal of Personality*, 69, 253–276.
- Mitchell, D. G. V., Colledge, E., Leonard, A., Blair, R. J. R. (2002). Risky decisions and response reversal: Is there evidence of orbito-frontal cortex dysfunction in psychopathic individuals? *Neuropsychologia*, 40, 2013-2022.
- Moffitt, T. E. (1993). Adolescence-limited and life-course-persistent anti-social behavior: A developmental taxonomy. *Psychological Review*, 100, 674-701.
- Morgan, A. P. & Lilienfeld, S. O. (2000). A meta-analytic review of the relation between antisocial behavior and neuropsychological measures of executive function. *Clinical Psychology Review*, 20, 113-156.
- Müller, J. L., Sommer, M., Wagner, V., Lamge, K., Taschler, H., Röder, H., Schuierer, G., Klein, H. E., & Hajak, G. (2003). Abnormalities in emotion processing within cortical and subcortical regions in criminal psychopaths: Evidence from a functional magnetic

- resonance imaging study using pictures with emotional content. *Biological Psychiatry*, *54*, 152-162.
- Newman, J. P. (1987). Reaction to punishment in extraverts and psychopaths: Implications for the impulsive behavior of disinhibited individuals. *Journal of Research in Personality*, *21*, 464-480.
- Newman, J. P., & Schmitt, W. A. (1998). Passive avoidance in psychopathic offenders: A replication and extension. *Journal of Abnormal Psychology*, *107*, 527-532.
- Newman, J. P., & Kosson, D. S. (1986). Passive avoidance learning in psychopathic and nonpsychopathic offenders. *Journal of Abnormal Psychology*, *95*, 252-256.
- Newman, J. P., & Lorenz, A. R. (2003). Response modulation and emotion processing: Implications for psychopathy and other dysregulatory psychopathology. In R. J. Davidson, K. Scherer, & H. H. Goldsmith (Eds.), *Handbook of Affective Sciences* (pp. 1043-1067). Oxford: Oxford University Press.
- Newman, J. P., Patterson, C. M., & Kosson, D. S. (1987). Response perseveration in psychopaths. *Journal of Abnormal Psychology*, *96*, 145-148.
- Newman, J. P., Schmitt, W. A., & Voss, W. D. (1997). The impact of motivationally neutral cues on psychopathic individuals: Assessing the generality of the response modulation hypothesis. *Journal of Abnormal Psychology*, *106*, 563-575.
- Ogloff, J. R., Wong, S. et Greenwood, A., (1990). Treating Criminal Psychopaths in a Therapeutic Community Program. *Behavioral Sciences and the Law*, *8*, 181-190.
- Patrick, C. J. (2001). Emotional processes in psychopathy. In A. Raine & J. Sanmartin (Eds.), *Violence and psychopathy* (pp. 57-77). New York: Kluwer Academic Publishers.

- Patrick, C. J. (2003, July). Psychopathy and Externalizing: Genetics and Brain Function, Paper presented at the *Developmental and Neuroscience Perspectives on Psychopathy* conference, Madison, Wisconsin.
- Patrick, C. J., Bradley, M. M., & Lang, P. J. (1993). Emotion in the criminal psychopath: Startle reflex modulation. *Journal of Abnormal Psychology, 102*, 82-92.
- Patrick, C. J., Curtin, J. J., & Tellegen, A. (2002). Development and validation of a brief form of the Multidimensional Personality Questionnaire (MPQ). *Psychological Assessment, 14*, 150-163.
- Patrick, C. J., & Zempolich, K. A. (1998). Emotion and aggression in the psychopathic personality. *Aggression and Violent Behavior, 3*, 303-338.
- Porter, S. & Woodworth, M. (2006), Psychopathy and Aggression. In C. Patrick (Ed.), *Handbook of Psychopathy* (pp. 481-494). New York: Guilford Press.
- Quinsey, V. L., Harris, G. T., Rice, M. E. & Cormier, C. A. (1998). *Violent offenders: Appraising and managing risk*. Washington, DC: American Psychological Association.
- Raine, A. (2001). Psychopathy, violence, and brain imaging. In A. Raine & J. Sanmartin (Eds.), *Violence and Psychopathy* (pp. 35-56). New York: Academic Press.
- Raine, A., Ishikawa, S. S., Arce, E., Lencz, T., Knuth, K. H., Birchle, S., LaCasse, L., & Coletti, P. (2004). Hippocampal structural asymmetry in unsuccessful psychopaths. *Biological Psychiatry, 55*, 185-191.
- Raine, A., Venables, P. H. & Williams, M. (1996). Better autonomic conditioning and faster electrodermal half-recovery time at age 15 years as possible protective factors against crime at age 29 years. *Developmental Psychology, 32*, 624-630.
- Riggins-Caspers, K. M., Cadoret, R. J. Knutson, J. F., & Langbehn, D. (2003). Biology-

- environment interaction and evocative, biology-environment correlation: Contributions of harsh discipline and parental psychopathology to problem adolescent behaviors, *Behavior Genetics*, 33, 205-220.
- Ruscio, J. (2000). Taxometric analysis with dichotomous indicators: The modified MAXCOV procedure and a case removal consistency test. *Psychological Reports*, 87, 929-939.
- Ruscio, J. (2006). *Estimating taxon base rates with the MAXCOV, MAXEIG, and MAMBAC taxometric procedures: A Monte Carlo study*. Manuscript in preparation.
- Ruscio, J. (2004, May). *Bootstrapping sampling distributions to quantify the fit of taxometric curves*. Paper presented at the annual meeting of the American Psychological Society, Chicago, IL.
- Ruscio, J., & Ruscio, A. M. (2004a). Clarifying boundary issues in psychopathology: The role of taxometrics in a comprehensive program of structural research. *Journal of Abnormal Psychology*, 113, 24-38.
- Ruscio, J., & Ruscio, A. M. (2004b). A conceptual and methodological checklist for conducting a taxometric investigation. *Behavior Therapy*, 35, 403-447.
- Ruscio, J., & Ruscio, A. M. (2004c). A nontechnical introduction to the taxometric method. *Understanding Statistics*, 3, 151-193.
- Ruscio, J., & Ruscio, A. M., & Keane, T. M. (2004). Using taxometric analysis to distinguish a small latent taxon from a latent dimension with positively skewed indicators: The case of Involuntary Defeat Syndrome. *Journal of Abnormal Psychology*, 113, 145-154.
- Ruscio, J., Ruscio, A. M., & Meron, M. (2005). *Applying the bootstrap to taxometric analysis: Generating empirical sampling distributions to help interpret results*. Manuscript submitted for publication.

- Salekin, R., T., Neumann, C. S., Leistico, A., M., DiCicco, T. M., & Duros, R, L. (in press). Construct validity of psychopathy in a youth offender sample: Taking a closer look at psychopathy's potential importance over disruptive behavior disorders. *Journal of Abnormal Psychology*.
- Schmauk, F. J., (1970). Punishment, arousal, and avoidance learning in sociopaths. *Journal of Abnormal Psychology*, 76, 325-335.
- Séguin, J. R. (2004). Neurocognitive elements of antisocial behaviour: Relevance of an orbitofrontal cortex account. *Brain and Cognition*, 55, 185-197.
- Skeem, J., Poythress, N., Edens, J., Lilienfeld, S., & Cale, E. (2003). Psychopathic personality or personalities? Exploring potential variants of psychopathy and their implications for risk assessment. *Aggression and Violent Behavior*, 8, 513-546.
- Skilling, T. A., Harris, G. T., Rice, M. E., & Quinsey, V. L. (2002). Identifying persistently antisocial offenders using the Hare Psychopathy Checklist and DSM Antisocial Personality Disorder criteria. *Psychological Assessment*, 14, 27-38.
- Skilling, T. A., Quinsey, V. L., Craig, W.M. (2001). Evidence of a taxon underlying serious antisocial behavior in boys. *Criminal Justice and Behavior*, 28, 450-470.
- Taylor, J., Loney, B. R., Bobadilla, L., Iacono, W. G., & McGue, M. (2003). Genetic and environmental influence on psychopathy: Findings from an adolescent male twin cohort. *Journal of Abnormal Child Psychology*, 31, 633-645.
- Tellegen, A., & Waller, N. G. (in press). *Exploring personality through test construction: Development of the multidimensional personality questionnaire*. Minneapolis, MN: University of Minnesota Press.

- Tiihonen, J., Hodgins, S., Vaurio, O., Laakso, M., Repo, E., Soininen, H., Aronen, H. J., Nieminen, P., & Savolainen, L. (2000). Amygdaloid volume loss in psychopathy. *Society for Neuroscience Abstracts*, 2017.
- Vasey, M. W., Kotov, R., Frick, P. J., & Loney, B. R. (2005). The latent structure of psychopathy in youth: A taxometric investigation. *Journal of Abnormal Child Psychology*, 33, 411-429.
- Viding, E., Blair, R. J. R., Moffitt, T. E., & Plomin, R. (in press). Evidence for substantial genetic risk for psychopathy in 7-year-olds. *Journal of Child Psychology and Psychiatry*.
- Vitacco, M. J., Rogers, R., Neumann, C. S., Harrison, K., & Vincent, G. (in press). A comparison of factor models on the PCL-R with mentally disordered offenders: The development of a four-factor model. *Criminal Justice and Behavior*.
- Vitacco, M. J., Neumann, C. S., & Jackson, R. (in press). Testing a four-factor model of psychopathy and its association with ethnicity, gender, intelligence, and violence. *Journal of Consulting and Clinical Psychology*.
- Völlm B., Richardson P., Stirling J., Elliott, R., Dolan, M., Chaudhry, I., del Ben, C., Mckie, S., Anderson, I., & Deakin, B. (2004). Neurobiological substrates of antisocial and borderline personality disorder: preliminary results of a functional fMRI study. *Criminal Behaviour and mental Health*, 14, 39-54.
- Waller, N. G. & Meehl, P. E. (1998). *Multivariate taxometric procedures: Distinguishing types from continua*. Newbury Park, CA: Sage.
- Welsh, G. S. (1956). Factor dimensions A and R. In G. S. Welsh & W. G. Dahlstrom (Eds.), *Basic readings on the MMPI in psychology and medicine* (pp. 264-281). Minneapolis: University of Minnesota Press.

Widiger, T. A. (1993). The DSM-III-R categorical personality disorder diagnoses: A critique and an alternative. *Psychological Inquiry*, 4, 75-90.

Widiger, T. A. & Costa, P. T. (1994). Personality and personality disorders. *Journal of Abnormal Psychology*, 103, 78-91.

Williamson, S. E., Harpur, T. J., & Hare, R. D. (1991). Abnormal processing of affective words by psychopaths. *Psychophysiology*, 28, 260-273.

Author note

Correspondence concerning this article should be addressed to Jean-Pierre Guay, School of Criminology, University of Montréal, C.P. 6128, Succ. Centre-ville, Montréal, Québec, Canada, H3C 3J7. Electronic mail may be sent to Jean-Pierre.Guay@Umontreal.ca

Table 1

Comparison Curve Fit Index (CCFI) Values for MAMBAC and MAXEIG Analyses

Indicator set	Full Sample	Max.	Med. Security (Samples 1, 5, 7, 8, 11, 13, & 14)	Min. Security (Samples 2, 3 & 6)	Blacks Only	Whites Only	Federal Sample (Sample 15)
		Security (Samples 9, 12 & 14)					
MAMBAC Analyses							
Summative	.272	.293	.325	.393	.279	.369	.318
Affective	.132	.420	.294	.332	.330	.294	.224
Antisocial	.420	.313	.402	.398	.361	.221	.400
Interpersonal	.220	.422	.247	.278	.414	.359	.337
Lifestyle	.429	.448	.471	.360	.356	.290	.201
PCL Factor 1	.407	.604	.362	.606	.566	.499	.485
PCL Factor 2	.500	.576	.472	.464	.460	.459	.447
MAXEIG Analyses							
Summative	.265	.338	.339	.422	.290	.345	.459
Affective	.291	.371	.431	.352	.359	.543	.423
Antisocial	.537	.501	.428	.426	.470	.403	.499
Interpersonal	.194	.395	.395	.379	.409	.376	.256
Lifestyle	.340	.421	.483	.480	.352	.477	.198
PCL Factor 1	.424	.438	.453	.503	.506	.565	.390
PCL Factor 2	.386	.508	.473	.408	.441	.486	.472

Note. CCFI values < .50 support dimensional structure and appear in bold print; CCFI values > .50 support taxonic structure.

Table 2

Summary of Taxon Base Rate Estimates

Indicator Set	MAMBAC		MAXEIG		L-Mode	
	# Estimates	<i>M</i> (<i>SD</i>)	# Estimates	<i>M</i> (<i>SD</i>)	Mode 1	Mode 2
Summative	4	.54 (.08)	4	.66 (.07)	.00	.80
Sim. Tax. Data		.65 (.07)		.43 (.02)		
Sim. Dim. Data		.54 (.04)		.62 (.06)		
Affective facet	4	.95 (.10)	6	.55 (.23)	—	—
Sim. Tax. Data		.74 (.15)		.59 (.16)		
Sim. Dim. Data		.94 (.12)		.50 (.20)		
Antisocial facet	5	.53 (.11)	10	.68 (.24)	—	—
Sim. Tax. Data		.56 (.05)		.61 (.15)		
Sim. Dim. Data		.51 (.07)		.56 (.20)		
Interpersonal facet	4	.56 (.21)	6	.42 (.11)	—	—
Sim. Tax. Data		.36 (.24)		.50 (.11)		
Sim. Dim. Data		.52 (.18)		.37 (.15)		
Lifestyle facet	5	.59 (.09)	10	.66 (.20)	—	—
Sim. Tax. Data		.57 (.05)		.63 (.09)		
Sim. Dim. Data		.55 (.08)		.61 (.13)		
PCL-R Factor 1	8	.60 (.11)	28	.52 (.18)	—	—
Sim. Tax. Data		.56 (.10)		.52 (.20)		
Sim. Dim. Data		.57 (.09)		.52 (.20)		
PCL-R Factor 2	10	.68 (.07)	45	.69 (.20)	—	—
Sim. Tax. Data		.60 (.06)		.60 (.17)		
Sim. Dim. Data		.69 (.14)		.61 (.18)		

Figure Captions


Figure 1. MAMBAC, MAXEIG, and L-Mode results for the four summative indicators. Within each 3-panel graph, the first panel shows the averaged curve for the research data and the second and third panels show the averaged curves for the simulated taxonic and dimensional comparison data. For comparison data, each sample's results are represented by a dotted line, with the average across all samples plotted as a solid line.

Figure 2. MAMBAC and MAXEIG results for indicator sets representing four theoretical facets of psychopathy. Aff. = affective; Anti. = antisocial; Int. = interpersonal; Life. = lifestyle. Within each 3-panel graph, the first panel shows the averaged curve for the research data and the second and third panels show the averaged curves for the simulated taxonic and dimensional comparison data. For comparison data, each sample's results are represented by a dotted line, with the average across all samples plotted as a solid line.


Figure 3. MAMBAC and MAXEIG results for indicator sets representing PCL-R Factors 1 and 2. Within each 3-panel graph, the first panel shows the averaged curve for the research data and the second and third panels show the averaged curves for the simulated taxonic and dimensional comparison data. For comparison data, each sample's results are represented by a dotted line, with the average across all samples plotted as a solid line.

Four Summative Indicators


MAMBAC



MAXEIG




L-Mode




MAMBAC

MAXEIG


Aff.




Anti.




Inter.




Life.




PCL Factor 1 – MAMBAC



PCL Factor 1 – MAXEIG



PCL Factor 2 – MAMBAC



PCL Factor 2 – MAXEIG

